

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/009263

International filing date: 21 March 2005 (21.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/555,380
Filing date: 22 March 2004 (22.03.2004)

Date of receipt at the International Bureau: 25 April 2005 (25.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1309564

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

April 18, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/555,380

FILING DATE: *March 22, 2004*

RELATED PCT APPLICATION NUMBER: *PCT/US05/09263*



Certified by

Don W. Duckas

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

22713 U.S. PTO
032204

ATTORNEY DOCKET NO. 19044.0057U1
Page 1 of 2

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a **PROVISIONAL APPLICATION FOR PATENT** under 37 C.F.R. § 1.53(c).

Docket Number		19044.0057U1		Type a Plus Sign (+) inside this box		+	
INVENTOR(s)							
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (City and Either State or Foreign Country)				
Subramaniam	Ananthan		1816 Forest Haven Lane, Birmingham, Alabama 35216				
TITLE OF INVENTION (500 characters max)							
NONPEPTIDE INHIBITORS OF MATRIX METALLOPROTEINASES							
CORRESPONDENCE ADDRESS							
Mitchell A. Katz Needle & Rosenberg, P.C. Customer Number 23859							
ENCLOSED APPLICATION PARTS (Check All That Apply)							
<input checked="" type="checkbox"/> Provisional Application Title Page <i>Number of Pages</i> [1]							
<input checked="" type="checkbox"/> Specification (includes Description, Claims, & Abstract) <i>Number of Pages</i> [57]							
<input checked="" type="checkbox"/> Drawing(s) <i>Number of Sheets</i> [1]							
<input checked="" type="checkbox"/> Authorization to Treat Reply Requesting Extension of Time as Incorporating Petition for Extension of Time [2 pages]							
<input checked="" type="checkbox"/> Other (specify): <u>Stamped, return postcard; Credit card payment form PTO-2038 (\$80).</u>							


17510 U.S. PTO
60555380
032204

METHOD PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (Check One)	
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR § 1.27.	FILING FEE AMOUNT \$ <u>80.00</u>
<input checked="" type="checkbox"/> A Credit Card Payment Form PTO-2038 is enclosed to cover the filing fees.	
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees.	
<input type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number _____.	
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any deficiency or credit any overpayment to Deposit Account No. <u>14-0629</u> .	

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.


- ☒ No.
- ☐ Yes. The name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

Signature  Date March 22, 2004

Typed or Printed Name: Mitchell A. Katz

Registration No. 33,919

CERTIFICATE OF EXPRESS MAILING UNDER 37 C.F.R. § 1.10	
I hereby certify that this correspondence and any items indicated as attached or included are being deposited with the United States Postal Service as Express Mail, Label No. EL 992019835 US, in an envelope addressed to: MAIL STOP PROVISIONAL PATENT APPLICATION, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450, on the date shown below.	
<u></u>	<u>3/22/04</u>
Chris C. Czajka	Date

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of)	
)	
Subramaniam Ananthan)	Art Unit: Unassigned
)	
Application No. Unassigned)	Examiner: Unassigned
)	
Filing Date: March 22, 2004)	Confirmation No. Unassigned
)	
For: NONPEPTIDE INHIBITORS OF)	
MATRIX METALLOPROTEINASES)	

AUTHORIZATION TO TREAT REPLY REQUIRING EXTENSION OF TIME
AS INCORPORATING PETITION FOR EXTENSION OF TIME

Mail Stop PROVISIONAL PATENT APPLICATION
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

NEEDLE & ROSENBERG, P.C.
Customer Number 23859

Sir:

Pursuant to 37 C.F.R. § 1.136(a)(3), the Commissioner is hereby requested and authorized to treat any concurrent or future reply in the above-identified application, requiring a petition for an extension of time for its timely submission, as incorporating a petition for extension of time for the appropriate length of time.


The Commissioner is hereby authorized to charge any fees which may be

ATTORNEY DOCKET NO. 19044.0057U1
PATENT


required, or credit any overpayment to Deposit Account No. 14-0629.

Respectfully submitted,

NEEDLE & ROSENBERG, P.C.


Mitchell A. Katz
Registration No. 33,919

NEEDLE & ROSENBERG, P.C.
Customer No. 23859

CERTIFICATE OF EXPRESS MAILING UNDER 37 C.F.R. § 1.10	
I hereby certify that this correspondence and any items indicated as attached or included are being deposited with the United States Postal Service as Express Mail, Label No. EL 992019835 US, in an envelope addressed to: Mail Stop PROVISIONAL PATENT APPLICATION , Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450, on the date shown below.	
 Chris Czajka	<u>3/22/04</u> Date

Express Mail No. EL 992019835 US
Attorney Docket No. 19044.0057U1
PATENT

PROVISIONAL APPLICATION
FOR
UNITED STATES LETTERS PATENT
FOR

NONPEPTIDE INHIBITORS OF MATRIX METALLOPROTEINASES

BY

**Subramaniam Ananthan, a citizen of the United States of America, residing at 1816
Forest Haven Lane, Birmingham, Alabama 35216.**

NONPEPTIDE INHIBITORS OF MATRIX METALLOPROTEINASES

BACKGROUND

Matrix metalloproteinases ("MMPs") are a class of zinc-dependent endopeptidase enzymes involved in the degradation and repair of major components of extracellular matrix and connective tissue. MMPs can be found in various cell types that reside in or are associated with connective tissues, such as fibroblasts, monocytes, macrophages, endothelial cells, and also invasive or metastatic tumor cells. MMPs are secreted from cells as latent proenzymes and are activated by Zn-dependent cleavage of the *N*-terminal part of the protein. When active MMPs are stimulated by growth factors and cytokines in the local tissue environment, they can degrade protein components of extracellular matrix and connective tissue, such as collagen, proteoglycans, fibronectin, and laminin. See H. Birkedal-Hansen, *Crit. Rev. Oral. Biol. Med.*, 1993, 4, 197-250.

Currently, it is known that there are fourteen different MMPs. These enzymes can be classified into several major categories according to their substrate specificities. For example, MMP-1, MMP-8, and MMP-13 are classified as collagenases. MMP-3 and MMP-11 are classified as stromelysins. MMP-2 and MMP-9 are classified as Type IV collagenases/gelatinases.

MMPs are of significant interest because they have been implicated in a wide variety of physiological and pathological conditions. Some examples of conditions known to be mediated by MMPs are tumor growth, osteoarthritis, rheumatoid arthritis, septic arthritis, restenosis, fibrosis, MMP-mediated osteopenias, inflammatory diseases of the central nervous system, reproduction, tissue morphogenesis, angiogenesis, skin aging, corneal ulceration, abnormal wound healing, bone disease, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the nervous system,

cirrhosis of the liver, glomerular disease of the kidney, premature rupture of fetal membranes, inflammatory bowel disease, periodontal disease, age related macular degeneration, diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, ocular inflammation, keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/neovascularization and corneal graft rejection. See M. Cockett, *et al.*, *Biochem. Soc. Symp.*, 1998, 63, 295-313; D. Keiner, *et al.*, *Can. Chemo. Pharm.*, 1999, 43, 42-51; D. Keiner, *Cancer Metastasis Rev.*, 1990, 9, 289-303; J. MacDougall, *et al.*, *Mol. Med. Today*, 2000, 64, 149-156; J. MacDougall, *et al.*, *Cancer Metastasis Rev.*, 1995, 14, 351-362; S. Curren, *et al.*, *Eur. J. Cancer*, 2000, 36, 1621-1630.

One particular area of research that has received much attention is the involvement of MMPs with cancer and the growth and spread of tumors. Indeed, the metastatic spread of cancer via proteolytic degradation of host biomatrix poses one of the greatest challenges in the treatment of cancer. Considerable evidence has been accumulated that indicates the involvement of MMPs in general, and of the gelatinases in particular, in local tumor growth, invasion, and metastatic spread of cancer to disseminated sites. For example, the level of expression of MMP-2 and MMP-9 is known to be elevated in certain tumor progression events. These enzymes degrade Type IV collagen, the major component of basement membranes, and denatured collagen (gelatin), leading to tumor metastasis. Also, the disruption of vascular membranes, composed mainly of Type IV collagen, by MMP-2 and MMP-9 is known to play a critical role in tumor metastasis.

Because of the involvement MMPs have in such a wide variety of physiological and pathological conditions, especially cancer and arthritis, synthetic inhibitors of these enzymes are considered attractive targets in drug discovery research. See J. B. Summers, *et al.*, *Ann. Rep. Med. Chem.*, 1998, 33, 131-140; A. H. Davidson, *et al.*, *Chem. Ind.*, 1997, 258-261; J. C. Spurlino, In "Structure-Based

Drug Design," Veerapandian, Ed., Marcel Dekker, Inc., N.Y., 1997, 171-189; R. P. Beckett, *et al.*, *Drug Disc. Today*, 1996, 1, 16-26. Such research pursuits have resulted in the development of several broad-spectrum peptidyl and partially selective nonpeptidyl MMP inhibitors as potential anticancer and antiarthritis agents.

5 See P. D. Brown, *Med. Oncology*, 1997, 14, 1-10; P. D. Brown, *APMIS*, 1999, 107, 174-180; P. D. Brown, *Expert Opin. Invest. Drugs*, 2000, 9, 2167-2177; J. Freskos, *et al.*, *Biorg. Med. Chem. Lett.*, 1999, 9, 943-948; L. J. MacPherson, *et al.*, *J. Med. Chem.*, 1997, 40, 2525-2532; M. Cheng, *et al.*, *J. Med. Chem.*, 2000, 43, 369-380. However, current results from both preclinical and clinical trials of MMP inhibitors

10 have been disappointing mainly due to poor bioavailability, poor selectivity, and undesirable side effects, such as tissue toxicity and even the promotion of liver metastasis. See "MMPs," Park W & Mecham R., AP, NY, 1998, pp. 1-14, 85-113, 115-149; M. Michaelides, *et al.*, *Curr. Pharma. Design*, 1999, 5, 787-819; E. Heath, *et al.*, *Drugs*, 2000, 59, 1043-1055; L. Seymour, *Cancer Treat. Rev.*, 1999, 25, 301-


15 312; K. Wocssner, *Ann. NY Aca. Sci.*, 1999, 878, 388-403; J. Skiles, *et al.*, *Ann. Rep. Med. Chem.*, 2000, 35, 167-176; M. Gowravaram, *et al.*, *J. Med. Chem.*, 1995, 38, 2570-2581; M. Gowravaram, *et al.*, *Biorg. Med. Chem. Lett.*, 1995, 5, 337-342; R. Greenwald, *et al.*, *Curr. Opin. Ther. Patents*, 1995, 4, 7-16; D. Levy, *et al.*, *J. Med. Chem.*, 1998, 41, 199-223; A. Kruger, *et al.*, *Cancer Res.*, 2001, 61, 1272-1275.

20 Therefore, in light of the clinical complexity associated with current MMP inhibitors, there is currently a need for new, potent inhibitors that more selectively target MMPs.

SUMMARY

In accordance with the purposes of the disclosed materials, compositions,

25 and methods, as embodied and broadly described herein, in one aspect, the disclosed subject matter relates to a compound having the following formula:



wherein X is $(\text{CH}_2)_n\text{O}$, $(\text{CH}_2)_n\text{S}$, $(\text{CH}_2)_n\text{NR}^1$, $(\text{CH}_2)_n(\text{CH}_2)$, or $\text{CH}=\text{CH}$, wherein $n = 0, 1$, or 2 ; R and R^1 are, independently, a substituted or unsubstituted alkyl, alkenyl, alkynyl, aryl, heteroaryl group, cycloalkyl, heterocycloalkyl, cycloalkenyl, or heterocycloalkenyl; and Z is NH or CH_2 ; or a pharmaceutically acceptable salt thereof.

In another aspect, the disclosed subject matter relates to a method for using the compounds described herein by administering an amount effective for modulation of matrix metalloprotease of at least one compound described herein to an environment comprising the matrix metalloprotease.

In yet another aspect, the disclosed subject matter relates to a method for using the compounds described herein by administering an amount effective for modulation of tumor metastasis of at least one compound described herein to a cell.

In a further aspect, the disclosed subject matter relates to a method for treating a subject with cancer comprising administering an effective amount of a compound described herein to a subject in need of the treatment.

In a still further aspect, the disclosed subject matter relates to a method for preventing cancer in a subject comprising administering an effective amount of a compound described herein to a subject.

In another aspect, the disclosed subject matter relates to a method for treating a subject with arthritis comprising administering an effective amount of a compound described herein to a subject in need of the treatment.

5 In still another aspect, the disclosed subject matter relates to selective modulators of matrix metalloproteinases and selective inhibitors of metalloproteinases. Also described herein are modulators of cancer metastasis and diseases such as arthritis. Further, methods of making and using such compounds are disclosed.

10 Additional advantages will be set forth in part in the description which follows, and in part will be obvious from the description, or may be learned by practice of the aspects described below. The advantages described below will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims. It is to be understood that both the foregoing general description and the following detailed description are exemplary and
15 explanatory only and are not restrictive.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawings, which are incorporated in and constitute a part of this specification, illustrate several aspects described below.

20 Figure 1A is a graph showing % inhibition of tumor invasion for various concentrations of compound 2c in the Amgel tumor invasion bioassay.

Figure 1B is a gelatin zymograph showing MMP-2 and MMP-9 activity inhibition with compound 2c.

DETAILED DESCRIPTION

The disclosed materials, compounds, compositions, and methods may be understood more readily by reference to the following detailed description of specific aspects of the materials and methods and the Examples included therein and to the Figure and the previous and following description.

5 Before the present materials, compounds, compositions, and/or methods are disclosed and described, it is to be understood that the aspects described below are not limited to specific synthetic methods or specific reagents, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular aspects only and is not intended to be limiting.

10 Disclosed are materials, compounds, compositions, and components that can be used for, can be used in conjunction with, can be used in preparation for, or are products of the disclosed method and compositions. These and other materials are disclosed herein, and it is understood that when combinations, subsets, interactions, groups, etc. of these materials are disclosed that while specific reference of each
15 various individual and collective combinations and permutation of these compounds may not be explicitly disclosed, each is specifically contemplated and described herein. For example, if a compound having a given formula is disclosed and discussed and a number of modifications that can be made to a number of R groups in the formula are discussed, each and every combination and permutation of the
20 compound and the modifications to the R groups that are possible are specifically contemplated unless specifically indicated to the contrary. Thus, if a class of substituents A, B, and C are disclosed as well as a class of substituents D, E, and F and an example of a combination molecule, A-D is disclosed, then even if each is not individually recited, each is individually and collectively contemplated. Thus, in
25 this example, each of the combinations A-E, A-F, B-D, B-E, B-F, C-D, C-E, and C-F are specifically contemplated and should be considered disclosed from disclosure of A, B, and C; D, E, and F; and the example combination A-D. Likewise, any

subset or combination of these is also specifically contemplated and disclosed. Thus, for example, the sub-group of A-E, B-F, and C-E are specifically contemplated and should be considered disclosed from disclosure of A, B, and C; D, E, and F; and the example combination A-D. This concept applies to all aspects of this disclosure including, but not limited to, steps in methods of making and using the disclosed compositions. Thus, if there are a variety of additional steps that can be performed it is understood that each of these additional steps can be performed with any specific embodiment or combination of embodiments of the disclosed methods, and that each such combination is specifically contemplated and should be considered disclosed.

Definitions

In this specification and in the claims which follow, reference will be made to a number of terms which shall be defined to have the following meanings:

As used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a compound" includes mixtures of compounds; reference to "an aryl substituent" includes mixtures of two or more such aryl substituents, and the like.

"Optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where the event or circumstance occurs and instances where it does not. For example, the phrase "optionally substituted aryl group" means that the aryl group may or may not be substituted and that the description includes both unsubstituted aryl groups and aryl groups where there is substitution.

Ranges may be expressed herein as from "about" one particular value and/or to "about" another particular value. When such a range is expressed, another aspect

includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent "about," it will be understood that the particular value forms another aspect. It will be further understood that the endpoints of each of the ranges are significant both in relation to
5 the other endpoint, and independently of the other endpoint.

References in the specification and concluding claims to parts by weight, of a particular element or component in a composition or article, denote the weight relationship between the element or component and any other elements or components in the composition or article for which a part by weight is expressed.
10 Thus, in a compound containing 2 parts by weight of component X and 5 parts by weight component Y, X and Y are present at a weight ratio of 2:5, and are present in such ratio regardless of whether additional components are contained in the compound.

A weight percent of a component, unless specifically stated to the contrary, is
15 based on the total weight of the formulation or composition in which the component is included.

The term "activity" as used herein refers to a biological activity. The term "pharmacological activity" as used herein refers to the inherent physical and/or chemical properties of a compound, molecule, modulator, or inhibitor. These
20 properties include but are not limited to efficacy, half-life, solubility, stability, affinity, and other pharmacokinetic and pharmacodynamic properties.

The terms "peptide" and "peptidyl" as used herein respectively refer to a class of compounds and chemical moieties composed of amino acids chemically bound together. In general, the amino acids are chemically bound together via amide
25 linkages (CONH). "Peptide" and "peptidyl" as used herein include oligomers of amino acids and small and large peptides, including polypeptides and proteins. The

terms "non-peptide" or "nonpeptidyl" refer to a class of compounds that are not composed of amino acids chemically bound together via an amide linkage.

As used herein, the term "substituted" is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, and aromatic and nonaromatic substituents of organic compounds. Illustrative substituents include, for example, those described below. The permissible substituents can be one or more and the same or different for appropriate organic compounds. For purposes of this disclosure, the heteroatoms, such as nitrogen, can have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valencies of the heteroatoms. This disclosure is not intended to be limited in any manner by the permissible substituents of organic compounds. Also, the terms "substitution" or "substituted with" include the implicit proviso that such substitution is in accordance with permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, *e.g.*, a compound that does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc.

The term "alkyl" as used herein is a branched or unbranched saturated hydrocarbon group of 1 to 24 carbon atoms, such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *t*-butyl, pentyl, hexyl, heptyl, octyl, decyl, tetradecyl, hexadecyl, eicosyl, tetracosyl and the like. The alkyl group can also be substituted or unsubstituted. The alkyl group can be substituted with one or more groups including, but not limited to, alkyl, halogenated alkyl, alkoxy, alkenyl, alkynyl, aryl, heteroaryl, aldehyde, amino, carboxylic acid, ester, halide, hydroxamate, hydroxy, ketone, nitro, silyl, sulfo-oxo, sulfonyl, sulfone, sulfoxide, or thiol, as described below. The term "halogenated alkyl" specifically refers to an alkyl group that is

substituted with one or more halide, *e.g.*, fluorine, chlorine, bromine, or iodine.

The term "alkoxy" as used herein is an alkyl group bound through a single, terminal ether linkage; that is, an "alkoxy" group may be defined as -OA where A is alkyl as defined above.

5 The term "alkenyl" as used herein is a hydrocarbon group of from 2 to 24 carbon atoms with a structural formula containing at least one carbon-carbon double bond. Asymmetric structures such as (AB)C=C(CD) are intended to include both the *E* and *Z* isomers. This may be presumed in structural formulae herein wherein an asymmetric alkene is present, or it may be explicitly indicated by the bond symbol
10 C=C.

The term "alkynyl" as used herein is a hydrocarbon group of 2 to 24 carbon atoms with a structural formula containing at least one carbon-carbon triple bond.

The term "aryl" as used herein is any carbon-based aromatic group including, but not limited to, benzene, naphthalene, phenyl, biphenyl, phenoxybenzene, etc.

15 The term "aromatic" also includes "heteroaryl," which is defined as an aromatic group that has at least one heteroatom incorporated within the ring of the aromatic group. Examples of heteroatoms include, but are not limited to, nitrogen, oxygen, sulfur, and phosphorus. The aryl group can be substituted or unsubstituted. The aryl group can be substituted with one or more groups including, but not limited to, alkyl,
20 halogenated alkyl, alkoxy, alkenyl, alkynyl, aryl, heteroaryl, aldehyde, amino, carboxylic acid, ester, halide, hydroxamate, hydroxy, ketone, nitro, silyl, sulfo-oxo, sulfonyl, sulfone, sulfoxide, or thiol as described herein. The term "biaryl" is a specific type of aryl group and is included in the definition of aryl. Biaryl refers to two aryl groups that are bound together via a fused ring structure, as in naphthalene,
25 or are attached via one or more carbon-carbon bonds, as in biphenyl.

The term "cycloalkyl" as used herein is a non-aromatic carbon-based ring

composed of at least three carbon atoms. Examples of cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc. The term "heterocycloalkyl" is a cycloalkyl group as defined above where at least one of the carbon atoms of the ring is substituted with a heteroatom such as, but not limited to, nitrogen, oxygen, sulfur, or phosphorus. The cycloalkyl group and heterocycloalkyl group can be substituted or unsubstituted. The cycloalkyl group and heterocycloalkyl group can be substituted with one or more groups including, but not limited to, alkyl, alkoxy, alkenyl, alkynyl, aryl, heteroaryl, aldehyde, amino, carboxylic acid, ester, halide, hydroxamate, hydroxy, ketone, nitro, silyl, sulfo-oxo, sulfonyl, sulfone, sulfoxide, or thiol as described herein.

The term "cycloalkenyl" as used herein is a non-aromatic carbon-based ring composed of at least three carbon atoms and contains at least one carbon-carbon double bond, C=C. Examples of cycloalkenyl groups include, but are not limited to, cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclopentadienyl, cyclohexenyl, etc. The term "heterocycloalkenyl" is a cycloalkenyl group as defined above where at least one of the carbon atoms of the ring is substituted with a heteroatom such as, but not limited to, nitrogen, oxygen, sulfur, or phosphorus. The cycloalkenyl group and heterocycloalkenyl group can be substituted or unsubstituted. The cycloalkenyl group and heterocycloalkenyl group can be substituted with one or more groups including, but not limited to, alkyl, alkoxy, alkenyl, alkynyl, aryl, heteroaryl, aldehyde, amino, carboxylic acid, ester, halide, hydroxamate, hydroxy, ketone, nitro, silyl, sulfo-oxo, sulfonyl, sulfone, sulfoxide, or thiol as described herein.

The term "aldehyde" as used herein is represented by the formula $-C(O)H$.

The terms "amine" or "amino" as used herein are represented by the formula NAA^1A^2 , where A, A^1 , and A^2 can be, independently, hydrogen, an alkyl, halogenated alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl group described above.

The term "carboxylic acid" as used herein is represented by the formula
-C(O)OH.

The term "ester" as used herein is represented by the formula -OC(O)A or
-C(O)OA, where A can be an alkyl, halogenated alkyl, alkenyl, alkynyl, aryl,
5 heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl group
described above.

The term "ether" as used herein is represented by the formula AOA¹, where
A and A¹ can be, independently, an alkyl, halogenated alkyl, alkenyl, alkynyl, aryl,
heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl group
10 described above.

The term "ketone" as used herein is represented by the formula AC(O)A¹,
where A and A¹ can be, independently, an alkyl, halogenated alkyl, alkenyl, alkynyl,
aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl
group described above.

15 The term "halide" as used herein refers to the halogens fluorine, chlorine,
bromine, and iodine.

The term "hydroxamate" as used herein is represented by the formula
-C(O)NHOH.

The term "hydroxyl" as used herein is represented by the formula -OH.

20 The term "nitro" as used herein is represented by the formula -NO₂.

The term "silyl" as used herein is represented by the formula -SiAA¹A²,
where A, A¹, and A² can be, independently, hydrogen, alkyl, halogenated alkyl,
alkoxy, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl,
or heterocycloalkenyl group described above.

The term "sulfo-oxo" as used herein is represented by the formulas $-S(O)A$, $-S(O)_2A$, $-OS(O)_2A$, or $-OS(O)_2OA$, where A can be hydrogen, an alkyl, halogenated alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl group described above.

5 The term "sulfonyl" is used herein to refer to the sulfo-oxo group represented by the formula $-S(O)_2A$, where A can be hydrogen, an alkyl, halogenated alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl group described above.

10 The term "sulfonamino" or "sulfonamide" as used herein is represented by the formula $-S(O)_2NH-$.

The term "sulfone" as used herein is represented by the formula $AS(O)_2A^1$, where A and A^1 can be, independently, an alkyl, halogenated alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl group described above.

15 The term "sulfoxide" as used herein is represented by the formula $AS(O)A^1$, where A and A^1 can be, independently, an alkyl, halogenated alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl group described above.

The term "thiol" as used herein is represented by the formula $-SH$.

20 "X," "Y," "R," " R^1 ," and " R^2 " as used herein can, independently, possess one or more of the groups listed above. For example, if R is a straight chain alkyl group, one of the hydrogen atoms of the alkyl group can optionally be substituted with a hydroxyl group, an alkoxy group, etc. Depending upon the groups that are selected, a first group can be incorporated within second group or, alternatively, the
25 first group can be pendant (*i.e.*, attached) to the second group. For example, with the

phrase "an alkyl group comprising a sulfonyl group," the sulfonyl group can be incorporated within the backbone of the alkyl group. Alternatively, the sulfonyl group can be attached to the backbone of the alkyl group. The nature of the group(s) that is (are) selected will determine if the first group is embedded or attached to the
5 second group.

As used herein, by a "subject" is meant an individual. Thus, the "subject" can include mammals (e.g., primate, human, etc.), domesticated animals (e.g., cats, dogs, etc.), livestock (e.g., cattle, horses, pigs, sheep, goats, etc.), laboratory animals (e.g., mouse, rabbit, rat, guinea pig, etc.), and birds. In one aspect, "subject" is a
10 mammal. In another aspect, "subject" is a human.

Reference to a "cell" herein can include a cell *in vitro*. Alternatively, reference to a "cell" can include a cell *in vivo*, which can be found in a subject. A "cell" can be a cell from any organism including, but not limited to, a bacterium, a eukaryote, or an animal.

15 By the term "effective amount" of a compound as provided herein is meant a nontoxic but sufficient amount of a compound to provide the desired result, e.g., modulation or inhibition. As will be pointed out below, the exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the disease that is being treated, the
20 particular compound used, its mode of administration, and the like. Thus, it is not possible to specify an exact "effective amount." However, an appropriate effective amount can be determined by one of ordinary skill in the art using only routine experimentation. Similarly, by the phrase "amount effective for modulation of a MMP" is meant a nontoxic but sufficient amount of a compound to modulate the
25 activity of at least one MMP. Also, by the phrase "amount effective for inhibition of a MMP" is meant a nontoxic but sufficient amount of a compound to inhibit the activity of at least one MMP. Again, the exact amount will vary from subject to

subject, depending on the species, age, and general condition of the subject, the severity of the disease that is being treated, the particular compound used, its mode of administration, and the like.


5 The phrase "environment comprising the MMP" is meant any environment where one or more MMP is present. Such environments can include, but are not limited to, subjects, organs, tumors, cells, gels, solutions, or neat MMP.

By "pharmaceutically acceptable" is meant a material that is not biologically or otherwise undesirable, *i.e.*, the material can be administered to an individual along with the selected compound without causing any undesirable biological effects or
10 interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained.

Reference will now be made in detail to specific aspects of the disclosed materials, compounds, compositions, components, and methods, examples of which are illustrated in the accompanying drawing.

15 Compounds

In one aspect, described herein are compounds having Formula I:



wherein X is $(CH_2)_nO$, $(CH_2)_nS$, $(CH_2)_nNR^1$, $(CH_2)_n(CH_2)$, or $CH=CH$, wherein $n = 0, 1$, or 2 ; R and R^1 are, independently, a substituted or unsubstituted alkyl, alkenyl, alkynyl, aryl, heteroaryl group, cycloalkyl, heterocycloalkyl, cycloalkenyl, or heterocycloalkenyl; and Z is NH or CH_2 ; or a pharmaceutically acceptable salt thereof.

In another aspect, described herein are compounds having Formula I, wherein Z is NH, R is a substituted or unsubstituted aryl or heteroaryl group; X is $(CH_2)_nO$, $(CH_2)_nS$, $(CH_2)_nNR^1$, $(CH_2)_n(CH_2)$, or $CH=CH$, wherein $n = 0, 1$, or 2 , and wherein R^1 is a substituted or unsubstituted alkyl, alkenyl, alkynyl, aryl, heteroaryl group, cycloalkyl, heterocycloalkyl, cycloalkenyl, or heterocycloalkenyl; or a pharmaceutically acceptable salt thereof.

In one aspect, described herein are compositions comprising a compound represented by Formula I.

The labels α and β are included in Formula I, as well as in other structures used herein, as aids to help identify and distinguish the particular carbon atom positions for further discussion. The choice of these labels is merely arbitrary and is not intended to be a limitation.

In the compounds represented by Formula I the pharmacophores, *e.g.*, the substituent that contains the sulfonyl group and the substituent that contains the hydroxamate group, are attached to adjacent carbon atoms, *i.e.*, carbons α and β . Also, the two carbon framework carrying the pharmacophores, *i.e.*, carbons α and β , is conformationally constrained by cyclic substitution. Further, in the pharmacophore containing the sulfonyl group, the sulfonyl group is positioned in the γ -position to the hydroxamate group, *i.e.*, there are three atoms in between the sulfonyl group and the hydroxamate group.


Compounds represented by Formula I can be optically active or racemic. The stereochemistry at carbons α and β can vary and will depend upon the spatial relationship between the substituent that contains the sulfonyl group and the hydroxamate group to one another. In one aspect, the stereochemistry at carbon α is S. In another aspect, the stereochemistry at carbon α is R. In one aspect, the stereochemistry at carbon β is S. In another aspect, the stereochemistry at carbon β is R. Using techniques known in the art, it is possible to vary the stereochemistry at carbons α and β .



Unless stated to the contrary, a formula with chemical bonds shown only as solid lines and not as wedges or dashed lines contemplates each possible isomer, e.g., each enantiomer and diastereomer, and a mixture of isomers, such as a racemic mixture.



Also described herein are the pharmaceutically acceptable salts of compounds represented by Formula I. Pharmaceutically acceptable salts are prepared by treating the hydroxamate and/or the sulfonamide with an appropriate amount of a pharmaceutically acceptable base. Representative pharmaceutically acceptable bases include ammonium hydroxide, sodium hydroxide, potassium hydroxide, lithium hydroxide, calcium hydroxide, magnesium hydroxide, ferrous hydroxide, zinc hydroxide, copper hydroxide, aluminum hydroxide, ferric hydroxide, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, lysine, arginine, histidine, and the like. In one aspect, the reaction is conducted in water, alone or in combination with an inert, water-miscible organic solvent, at a temperature of from about 0°C to about 100°C, such as at room temperature. The molar ratio of compounds represented by Formula I to be used is chosen to provide the ratio desired for any particular salts. For preparing, for example, the ammonium salts of

the hydroxamate, the hydroxamate can be treated with approximately one equivalent of pharmaceutically acceptable base to yield a neutral salt.

In one aspect, the R group in Formula I can be a substituted aryl group that is represented by the following formula.



wherein R^2 is Br; methoxy; ; , wherein Y = O, S, or CH₂;







5

In another aspect, the R group in Formula I can be *p*-methoxyphenyl, *p*-biphenyl, *p*-phenoxyphenyl, *p*-(phenylethynyl)phenyl, *p*-(phenylethenyl)phenyl, and linear tricyclic systems with three aryl groups or two aryl groups tethered to a central piperidiny ring system and their heteroaromatic analogues.

10

A non-exhaustive list of specific examples of compounds that are represented by Formula I are shown below.



a: $R^2 = \text{Br}$

b: $R^2 = \text{methoxy}$

c: $R^2 =$

d: $R^2 =$

e: $R^2 = (\text{CH}_3)_2\text{N}-$

f: $R^2 =$

g: $R^2 =$

h: $R^2 =$

i: $R^2 =$


j: $R^2 =$


In the list above, any of the R^2 substituents on compounds 1 through 8 can be any of the R^2 substituents labeled a through j. Also, specific compounds are referred to herein by listing the number of the formula above (e.g., 1 through 8) along with

the letter of the R^2 substituent (e.g., a through j). For example, a compound of formula 1 with R^2 as Br (i.e., R^2 is the substituent labeled "a") can be referred to as compound "1a."

In the list above, compounds 1-4 are considered sulfonamides and compounds 5-8 are considered sulfones. Among the sulfonamides, compounds 1 and 2 possess a saturated, cyclohexane framework, whereas compounds 3 and 4 possess an unsaturated, cyclohexene framework. Similarly, among the sulfones, compounds 5 and 6 possess a saturated, cyclohexane framework while compounds 7 and 8 possess an unsaturated, cyclohexene framework.


On a cyclic framework containing at least two adjacent substituents, as is shown in Formula I, there can be two chiral centers depending on the substituents. Also, the relative stereochemical orientation of the two substituents can be either *cis* or *trans*. Thus, compounds 1, 3, 5, and 7 represent *cis* isomers while compounds 2, 4, 6, and 8 represent *trans* isomers. Each of the *cis* or *trans* structures represents a racemate consisting of two enantiomers of opposite absolute configurations. For example, the *cis*-biphenylsulfonamide (1c) represents two *cis* isomers, one with an $\alpha S, \beta R$ configuration and the other with an $\alpha R, \beta S$ configuration. Similarly, the *trans*-biphenylsulfonamide (2c) represents two *trans* isomers, one with an $\alpha S, \beta S$ configuration and the other with an $\alpha R, \beta R$ configuration. The structures of these particular compounds, 1c and 2c, are shown below.






(2c)


Other specific examples include,



(e.g., compound 3-4c),



(e.g., compound 5-6d), and



As described herein, compounds represented by Formula I are modulators of MMPs. While not wishing to be bound by theory, it is believed that since compounds represented by Formula I have hydroxamate groups that can bind Zn, compounds represented by Formula I can be modulators of all Zn-dependent MMPs, for example, MMP-1, MMP-3, MMP-7, MMP-8, MMP-11, and MMP-13, or a

mixture thereof. In this respect, compounds represented by Formula I can be said to be broad spectrum MMP modulators.

In another aspect, compounds represented by Formula I are selective modulators of MMPs. In still another aspect, compounds represented by Formula I
5 are selective modulators of MMP-2 and MMP-9. In yet another aspect, compounds represented by Formula I are capable of modulating tumor progression, tumor metastasis, and tumor invasion, and are also antiarthritic agents.

In another aspect, compounds represented by Formula I are potent inhibitors of MMPs. While not wishing to be bound by theory, it is believed that since
10 compounds represented by Formula I have hydroxamate groups that can bind Zn, compounds represented by Formula I can be inhibitors of all Zn-dependent MMPs, for example, MMP-1, MMP-3, MMP-7, MMP-8, MMP-11, and MMP-13, or a mixture thereof. In this respect, compounds represented by Formula I can be said to be broad spectrum MMP inhibitors.

15 In one aspect, compounds represented by Formula I are selective inhibitors of MMPs. In another aspect, compounds represented by Formula I are selective inhibitors of MMP-2 and MMP-9. In yet another aspect, compounds represented by Formula I are capable of inhibiting tumor progression, tumor metastasis, and tumor invasion, and are also antiarthritic agents.


20 **Synthetic Methods**

Compounds represented by Formula I can be readily synthesized using techniques generally known to those of skill in the art. The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Co., (Milwaukee, Wis.), Acros Organics (Morris
25 Plains, N.J.), Fisher Scientific (Pittsburgh, Pa.), or Sigma (St. Louis, Mo.) or are prepared by methods known to those skilled in the art following procedures set forth

in references such as Fieser and Fieser's Reagents for Organic Synthesis, Volumes 1-17 (John Wiley and Sons, 1991); Rodd's Chemistry of Carbon Compounds, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989); Organic Reactions, Volumes 1-40 (John Wiley and Sons, 1991); March's Advanced Organic
5 Chemistry, (John Wiley and Sons, 4th Edition); and Larock's Comprehensive Organic Transformations (VCH Publishers Inc., 1989).

In one aspect, compounds represented by Formula I can be prepared by methods illustrated in Schemes I and II. These schemes are merely illustrative of some methods by which the compounds disclosed herein can be synthesized, and
10 various modifications to these schemes can be made and will be apparent to one skilled in the art having reviewed this disclosure.

In the following discussion, the starting materials and the intermediates of the reactions can be isolated and purified, if desired, using conventional techniques, including but not limited to, filtration, distillation, crystallization, chromatography,
15 and the like. Such materials can be characterized using conventional means, including physical constants and spectral data. Also, unless specified to the contrary, the reactions described herein can take place at atmospheric pressure over a temperature range from about -78°C to about 150°C, from about 0°C to about 125°C, or at about room (or ambient) temperature, e.g., about 20°C.



Scheme I provides an outline of a synthetic route for accessing the sulfonamide compounds represented by Formula I, wherein $Z = NH$, e.g., compounds 1, 2, 3, and 4 discussed above, starting with racemic *cis*- or *trans*-cyclic amino acid starting material (9). Starting material 9 is commercially available or is synthetically accessible by methods known to those skilled in the art. For example, starting materials *cis*-2-aminocyclohexanecarboxylic acid (racemic) or *trans*-2-aminocyclohexanecarboxylic acid (racemic), which lead to compounds 1 and 2, respectively, are available from commercial suppliers such as Acros Organics (Morris Plains, N.J.). Similarly, the use of cyclohexene analogues of 9 as starting materials provide the corresponding cyclohexene compounds 3 and 4. To arrive at individual enantiomers of the final compounds, known chiral, enantiomerically pure cyclic amino acid analogs of 9 can be used as the starting materials. Synthetic procedures for the preparation of enantiomerically pure cyclic amino acids are also known in the art. See N. Harmat, et. al., *Bioorg. Med. Chem. Lett.*, 1998, 8, 1249-

1254, which is incorporated by reference herein for its teachings of synthetic procedures for the preparation of enantiomerically pure cyclic amino acids.


In Scheme I, starting material 9 is reacted with a functional derivative of sulfonic acid, $R-SO_2LG$, where LG represents a suitable leaving group, such as, a chloride, anhydride, or mixed anhydride. This reaction can take place under basic conditions suitable to provide the sulfonamide. Suitable bases for this reaction are well known and include, but are not limited to, carbonates, bicarbonates, hydroxides, alkoxides, hydrides, and amines, such as trimethylamine, triethylamine, diisopropylamine, *N*-ethyl-diisopropyl amine, pyridine, or dimethylaminopyridine, including a mixture thereof. Also, the reaction can be carried out in the presence of an organic solvent, such as dioxane, dichloromethane, 1,2-dichloroethane, 1,1,1-trichloroethane, *N,N*-dimethylformamide (DMF), *N,N*-dimethylacetamide, dimethylsulfoxide (DMSO), acetonitrile, ethyl acetate, ether, benzene, toluene, or xylene, including a mixture thereof. In one aspect, the reaction can be carried out with starting material 9 and a sulfonyl chloride ($R-SO_2Cl$) in the presence of sodium carbonate in dioxane-water solvent.

The resulting sulfonamide is subsequently coupled with a protected hydroxylamine, H_2N-OPG , where PG represents a protecting group, under suitable amino-acid coupling conditions. Alternatively, the acid function in the sulfonamide is activated, for example, by conversion to the acid chloride or mixed anhydride, and then reacted with a protected hydroxylamine. Protected hydroxyl amines are commercially available or can be prepared by methods known in the art. Typically, protected hydroxylamines are prepared by reacting hydroxylamine with a suitable protecting group. The protecting group that is used will depend on the specific reaction conditions, other substituents that may be present, availability, or preference.

Conditions for coupling the protected hydroxylamine and the sulfonamide are well known in the art and typically involve contacting the sulfonamide with the protected hydroxylamine in the presence of one or more activating agents. Various activating agents that can be used for the coupling reaction include, but are not limited to, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC), dicyclohexylcarbodiimide (DCC), *N,N*-diisopropyl-carbodiimide (DIP), benzotriazol-1-yl-oxy-tris-(dimethylamino)phosphonium hexa-fluorophosphate (BOP), hydroxybenzotriazole (HOBt), and *N*-methylmorpholine (NMM), including a mixture thereof. The coupling reaction can be carried out in *N*-methylpyrrolidone (NMP) or in DMF. In one aspect, the coupling reaction can involve the treatment of the sulfonamide with a protected hydroxylamine in the presence of EDC, HOBt, and NMM in DMF. See Y. Tamura, *et al.*, *J. Med. Chem.*, 1998, 41, 640-649, which is incorporated by reference herein for its teaching of amine-acid coupling reactions.

Conditions for converting acid function to a reactive derivative such as an acid chloride, for example using thionyl chloride or oxalyl chloride, or to an anhydride, for example by reaction with chloroformic esters under appropriate conditions, followed by reaction of these activated intermediates with or without isolation with hydroxylamine or a protected hydroxylamine are known in the art and can be applied as an alternative method to the coupling of the acid with a protected hydroxylamine. See Y. Tamura, *et al.*, *J. Med. Chem.*, 1998, 41, 640-649.; P. O'Brien, *et al.*, *J. Med. Chem.*, 2000, 43, 156-166; M. Gowravaram, *et al.*, *J. Med. Chem.*, 1995, 38, 2570-2581, which are incorporated herein by reference for their teachings of the preparation and reactions of activated acids.

The final step of Scheme I involves the removal of the protecting group PG under hydrolytic conditions to result in compounds represented by Formula I, wherein Z = NH. Suitable conditions for the removal of the protecting group are discussed later.



Scheme II provides an outline of a synthetic route for accessing the sulfone compounds represented by Formula I, wherein $Z = \text{CH}_2$, e.g., compounds 5, 6, 7, and 8 discussed above, starting with racemic *cis*- or *trans*-lactone (10). Scheme II is based on the ring opening reaction of lactones with thiols (R-SH) in the presence of Lewis acid catalysts. Starting material 10 is commercially available or can be synthesized by methods known in the art. To arrive at individual enantiomers of the final compounds, known chiral, enantiomerically pure lactone analogs of 10 can be used as the starting materials. Synthetic procedures for the preparation of enantiomerically pure lactones are also known in the art. See D. Bailey, *et al.*, *J. Org. Chem.*, 1970, 35, 3574-3576; P. Kennewell, *et al.*, *J. Chem. Soc. Perkin. Trans. I*, 1982, 2563-2570, which are incorporated by reference herein for their teachings of synthetic procedures for the preparation of enantiomerically pure lactones.

The use of appropriate cyclohexane analogs of starting material 10 provide corresponding compounds 5 and 6, whereas the use of appropriate cyclohexene

analogs of starting material 10 provide corresponding compounds 7 and 8. The thiol R-SH is commercially available or can be synthesized by methods known in the art.

Lewis acids that are suitable for the ring opening reaction of the lactone 10 are well known in the art. For example, suitable Lewis acids include, but are not limited to, AlCl_3 , AlBr_3 , SO_3 and complexes of SO_3 , BF_3 , BF_3 etherate, ZnCl_2 , TiCl_4 , SbF_5 , SnCl_4 and the like, including a mixture thereof. Suitable solvents include, for example, dichloromethane, 1,2-dichloroethane, 1,1,1-trichloroethane, *N,N*-dimethylformamide (DMF), *N,N*-dimethylacetamide, dimethylsulfoxide (DMSO), acetonitrile, ethyl acetate, ether, benzene, toluene, or xylene, including a mixture thereof.

After the lactone is opened, the sulfur is oxidized by methods known in the art. Various oxidizing agents and conditions are discussed in Hudlicky, *Oxidations in Organic Chemistry*, ACS monograph 186, 1990, which is incorporated by reference herein for its teachings of oxidation reactions. Suitable oxidation agents include, for example, hydrogen peroxide, sodium meta-periodate, oxone (potassium peroxy monosulfate), meta-chloroperoxybenzoic acid, periodic acid and the like, including a mixture thereof. Suitable solvents include, for example, acetic acid (for sodium meta-periodate) and, for other peracids, ethers such as THF and dioxane, and acetonitrile, DMF and the like, including a mixture thereof. For oxone, suitable solvents include, for example, aqueous alcohols, such as methanol, ethanol, and propanol, including a mixture thereof.

Finally, in Scheme II, a protected hydroxylamine ($\text{H}_2\text{N}-\text{OPG}$) is coupled to the sulfone and the protecting group PG removed to result in compounds represented by Formula I. The protected hydroxylamine, as mentioned above, is commercially available or can be prepared by synthetic methods known in the art. Also, the conditions for the coupling reactions are analogous to those coupling conditions discussed above in Scheme I.

The phrase "protecting group" as used herein means a chemical moiety that temporarily modifies a potentially reactive functional group and protects the functional group from undesired chemical transformations. Protecting group chemistry is known to one of skill in the art. See T. Greene, *et al.*, "Protective Groups in Organic Synthesis," 2nd ed., Wiley, N.Y., 1991, which is incorporated by
5 reference herein for its teaching of protecting groups and methods of adding and removing protecting groups.

Examples of protecting groups suitable for use with the protected hydroxylamine in either Scheme I or II include, but are not limited to, tertbutyl,
10 benzyl, tetrahydropyranyl, and silyl ethers such as trimethylsilyl, tert-butyl dimethylsilyl, and triisopropylsilyl. Hydrolytic and hydrogenolytic conditions for removing protecting groups used herein, which thus reveals the hydroxamate group, are generally well known and will depend on the particular protecting group used. Hydrolytic deprotection can generally be accomplished by hydrolysis under
15 basic conditions or in the presence of a suitable acid, such as hydrochloric acid, hydrobromic acid, acetic acid, trifluoroacetic acid, or by contact with a suitable acidic resin, such as AMBERLYSTTM (Rohm Haas, Philadelphia, PA) or DOWEXTM (Dow, Midland, Mich.). Suitable solvents include, for example, dichloromethane, 1,2-dichloroethane, 1,1,1-trichloroethane, *N,N*-dimethylformamide
20 (DMF), *N,N*-dimethylacetamide, dimethylsulfoxide (DMSO), acetonitrile, ethyl acetate, ether, benzene, toluene, or xylene, including a mixture thereof.

Hydrogenolytic removal of a protecting group such as a benzyl group can be carried out by hydrogenation in the presence of a catalyst such as palladium on carbon at an appropriate temperature and pressure in a suitable solvent.

25 While the synthetic routes discussed above can be performed as solution-phase multiple parallel syntheses, which involves the synthesis of compounds in individual reaction vessels, other methods can be performed. For example,

combinatorial based syntheses or solid phase syntheses can be used and will depend on the particular compounds to be synthesized, the availability of reagents, or preference.

Utility and Administration

5 The compounds represented by Formula I have many uses. For example, the compounds have uses in areas where modulation or inhibition of MMPs is therapeutically or prophylactically beneficial, such as in treating or preventing cancer and arthritis. In one aspect the compounds represented by Formula I are used to treat a subject with cancer. Such cancer can include, but is not limited to, carcinoma,
10 melanoma, leukemia, or adenoma.

As noted above, MMPs are suitable targets for anticancer agents because Type IV collagenolytic activity is required by metastasizing tumor cells to traverse tissue/vascular membrane barriers. Also, over-production of MMPs clinically correlates with invasive and metastatic behavior of various tumors. Therefore,
15 administering an effective amount of a compound represented by Formula I to modulate or inhibit MMPs, such as MMP-2 and/or MMP-9, can modulate or inhibit such tumor progression events as tumor metastasis, tumor invasion, and angiogenesis.

In one aspect, the compounds represented by Formula I can selectively
20 modulate or inhibit various MMPs. Recently, information regarding the three-dimensional structure of the catalytic domain of several MMPs, e.g., MMP-1, -3, -7, and -8, and of the catalytic domain-inhibitor complexes, determined by X-ray crystallography and NMR-spectroscopy, has become available. See Y. Tamura, *et al.*, *J. Med. Chem.*, 1998, 21, 640-649; R. Kiyama, *et al.*, *J. Med. Chem.*, 1999, 42,
25 1723-1738; C. J. Burns, *et al.*, *Angew. Chem. Int. Ed.*, 1998, 37, 2848-2850; L. E. Burgess, *et al.*, *Chem. Abst.*, 1998, 128, 127820; A. Pavlovsky, *et al.*, *Protein Sci.*,

1999, 8, 1455-1462; B. Lovejoy, *et al.*, *Science*, 1994, 263, 375-377; T. Stams, *et al.*,
Nature: Struct. Biol., 1994, 1, 119-123; W. Bode, *et al.*, *EMBO J.*, 1994, 13, 1263;
B. Stockman, *et al.*, *Protein Sci.*, 1998, 7, 2118-2126; 2281-2286. These studies
indicate that the core structure of MMPs consists of three alpha helices and five
5 stranded beta sheet and a catalytic zinc ion located at the bottom of the catalytic
cleft, coordinated by three histidine ("His") residues. One difference, however,
among the various MMPs, is in the shape and size of the S1' pocket, an additional
substrate binding domain embedded within the catalyst Zn-binding domain. In
contrast to the open S1, S2', and S3' subsites, the S1' pocket penetrates into the core
10 of the MMP enzyme. This S1' pocket is relatively shallow in MMP-1 and MMP-7,
and narrow in MMP-9, while being a much deeper channel in MMP-2, MMP-3, and
MMP-8.

While not wishing to be bound by theory, it is believed that this unique
conformation of the S1' activity domains in MMPs, such as MMP-2 and MMP-9,
15 contributes to the specificity of the disclosed compound. Compounds represented by
Formula I, which incorporate conformational restraint at the α - and β -position of the
hydroxamate group with a sulfonyl substituent extending from the exocyclic γ -
position, can selectively occupy the deep S1' pocket of the active sites of MMP-2
and MMP-9. Also, while not wishing to be bound by theory, it is believed that the
20 conformationally restrained framework present in compounds represented by
Formula I aids in the projection of the sulfonyl substituent toward the S1' pocket,
whereas lack of conformational restraint is believed to lead to inhibitors lacking
potency due to greatly increased conformational flexibility.

In one aspect, a method for using compounds represented by Formula I
25 comprises administering an amount effective for modulation of a MMP of at least
one compound represented by Formula I to an environment comprising the MMP.

The MMP can be any MMP or mixtures of MMPs. In one aspect, the MMP is MMP-2, MMP-9, or a mixture thereof.

5 In another aspect, a method for using compounds represented by Formula I comprises administering an amount effective for inhibition of a MMP of at least one compound represented by Formula I to an environment comprising the MMP. The MMP can be any MMP or mixtures of MMPs. In one aspect, the MMP is MMP-2, MMP-9, or a mixture thereof.

10 In yet another aspect, an amount effective for modulation of a MMP of at least one compound represented by Formula I is administered to an environment comprising the MMP. In still another aspect, an amount effective for inhibition of a MMP of at least one compound represented by Formula I is administered to an environment comprising the MMP. Again, the MMP can be any MMP, *e.g.*, MMP-2, MMP-9, or mixtures of MMPs. In one aspect, the compound represented by Formula I that is administered to an environment comprising a MMP is compound
15 1c, 2c, or a mixture thereof.

The administration of compounds represented by Formula I to an environment comprising a MMP can be conducted *in vivo* or *in vitro*.

20 In one aspect, a method for using a compound represented by Formula I comprises administering an amount effective for modulation of tumor metastasis of at least one compound represented by Formula I to a subject or cell. In another aspect, a method for using a compound represented by Formula I comprises administering an amount effective for inhibition of tumor metastasis of at least one compound represented by Formula I to a subject or cell. In still another aspect, the cell is a HT-1080 cell.

25 In one aspect, the amount effective for modulation is equivalent to an amount effective for inhibition. Modulation and/or inhibition of the MMP can be measured

by methods known in the art. In one aspect, modulation can be measured by any change in tumor invasion and inhibition can be measured by arrest of tumor invasion. In another aspect, modulation can be measured by any change in tumor angiogenesis and inhibition can be measured by arrest of tumor angiogenesis.

- 5 Inhibition potency can be characterized by an IC_{50} less than about 3000nM, less than about 1500nM, less than about 1000nM, less than about 500nM, or less than about 200nM.

In one aspect, the compounds described herein can be administered to a subject in need of treatment, such as a subject with cancer or arthritis. The subject in
10 need of treatment can comprise a human or an animal including, but not limited to, a rodent, dog, cat, horse, bovine, ovine, or non-human primate and the like, that is in need of alleviation or amelioration from a recognized medical condition.

Any of the compounds represented by Formula I can be delivered to a subject as part of a cocktail, *i.e.*, used in combination with other pharmaceutical agents. For
15 example, the compounds represented by Formula I can be part of an anti-cancer cocktail, *i.e.*, one or more of the compounds represented by Formula I can be used with one or more anti-cancer agents. The use of cocktails in the treatment of cancer is routine. In one aspect, an anti-cancer cocktail involves a common administration vehicle (*e.g.*, pill, tablet, implant, injectable solution, etc.) that can contain effective
20 amounts of both one or more compounds represented by Formula I and one or more anti-cancer drugs. Alternatively, an anti-cancer cocktail involves the sequential, simultaneous, or scheduled administration of effective amounts of one or more compounds represented by Formula I and one or more anti-cancer drugs. For example, one or more compounds represented by Formula I can first be given to a
25 subject and then, after some period of time, an anti-cancer agent is given to the subject. The order of administration can be determined by one skilled in the art

depending on factors such as the particular type of anti-cancer drug, the type and severity of the cancer, and the like.

Suitable anti-cancer agents that can be used in an anti-cancer cocktail with the compounds represented by Formula I include, but are not limited to, Acivicin;
5 Aclarubicin; Acodazole Hydrochloride; AcrQnine; Adozelesin; Aldesleukin; Altretamine; Ambomycin; Ametantrone Acetate; Aminoglutethimide; Amsacrine; Anastrozole; Anthramycin; Asparaginase; Asperlin; Azacitidine; Azetepa; Azotomycin; Batimastat; Benzodepa; Bicalutamide; Bisantrone Hydrochloride; Bisnafide Dimesylate; Bizelesin; Bleomycin Sulfate; Brequinar Sodium;
10 Bropirimine; Busulfan; Cactinomycin; Calusterone; Caracemide; Carbetimer; Carboplatin; Carmustine; Carubicin Hydrochloride; Carzelesin; Cedefingol; Chlorambucil; Cirolemycin; Cisplatin; Cladribine; Crisnatol Mesylate; Cyclophosphamide; Cytarabine; Dacarbazine; Dactinomycin; Daunorubicin Hydrochloride; Decitabine; Dexormaplatin; Dezaguanine; Dezaguanine Mesylate;
15 Diaziquone; Docetaxel; Doxorubicin; Doxorubicin Hydrochloride; Droloxifene; Droloxifene Citrate; Dromostanolone Propionate; Duazomycin; Edatrexate; Eflomithine Hydrochloride; Elsamitrucin; Enloplatin; Enpromate; Epipropidine; Epirubicin Hydrochloride; Erbulozole; Esorubicin Hydrochloride; Estramustine; Estramustine Phosphate Sodium; Etanidazole; Ethiodized Oil I 131; Etoposide;
20 Etoposide Phosphate; Etoprine; Fadrozole Hydrochloride; Fazarabine; Fenretinide; Floxuridine; Fludarabine Phosphate; Fluorouracil; Flurocitabine; Fosquidone; Fostriecin Sodium; Gemcitabine; Gemcitabine Hydrochloride; Gold Au 198; Hydroxyurea; Idarubicin Hydrochloride; Ifosfamide; Ilmofosine; Interferon Alfa-2a; Interferon Alfa-2b; Interferon Alfa-n1; Interferon Alfa-n3; Interferon Beta- I a;
25 Interferon Gamma- I b; Iproplatin; Irinotecan Hydrochloride; Lanreotide Acetate; Letrozole; Leuprolide Acetate; Liarozole Hydrochloride; Lometrexol Sodium; Lomustine; Losoxantrone Hydrochloride; Masoprocil; Maytansine; Mechlorethamine Hydrochloride; Megestrol Acetate; Melengestrol Acetate;

Melphalan; Menogaril; Mercaptopurine; Methotrexate; Methotrexate Sodium;
Metoprine; Meturedapa; Mitindomide; Mitocarcin; Mitocromin; Mitogillin;
Mitomalcin; Mitomycin; Mitosper; Mitotane; Mitoxantrone Hydrochloride;
Mycophenolic Acid; Nocodazole; Nogalamycin; Ormaplatin; Oxisuran; Paclitaxel;
5 Pegaspargase; Peliomycin; Pentamustine; Peplomycin Sulfate; Perfosfamide;
Pipobroman; Pipo sulfan; Piroxantrone Hydrochloride; Plicamycin; Plomestane;
Porfimer Sodium; Porfiromycin; Prednimustine; Procarbazine Hydrochloride;
Puromycin; Puromycin Hydrochloride; Pyrazofurin; Riboprine; Rogletimide;
Safingol; Safingol Hydrochloride; Semustine; Simtrazene; Sparfosate Sodium;
10 Sparsomycin; Spirogermanium Hydrochloride; Spiromustine; Spiroplatin;
Streptonigrin; Streptozocin; Strontium Chloride Sr 89; Sulofenur; Talisomycin;
Taxane; Taxoid; Taxol; Tecogalan Sodium; Tegafur; Teloxantrone Hydrochloride;
Temoporfin; Teniposide; Teroxirone; Testolactone; Thiamiprine; Thioguanine;
Thiotepa; Tiazofurin; Tirapazamine; Topotecan Hydrochloride; Toremifene Citrate;
15 Trestolone Acetate; Triciribine Phosphate; Trimetrexate; Trimetrexate Glucuronate;
Triptorelin; Tubulozole Hydrochloride; Uracil Mustard; Uredapa; Vapreotide;
Verteporfin; Vinblastine Sulfate; Vincristine Sulfate; Vindesine; Vindesine Sulfate;
Vinepidine Sulfate; Vinglycinate Sulfate; Vinleurosine Sulfate; Vinorelbine Tartrate;
Vinrosidine Sulfate; Vinzolidine Sulfate; Vorozole; Zeniplatin; Zinostatin; and
20 Zorubicin Hydrochloride.

Modulators or inhibitors of MMPs can provide synergistic effects when used
in a cocktail. For example, the use of other MMP modulators or inhibitors in
combination with anti-cancer agents has been known to produce synergistic effects,
i.e., the amount of modulation or inhibition observed when using a combination of
25 MMP modulator or inhibitor and anti-cancer agent is greater than the amount of
modulation or inhibition observed when either is used alone. *See M. Ohta et al.,*
Japanese J. Cancer. Res., 2001, 92, 688; *M. Maki et al., Clin. Exp. Metastasis,*

2002, 19, 519. Accordingly, the use of compounds represented by Formula I can provide synergistic effects when administered as part of an anti-cancer cocktail.

The dosages or amounts of the compounds described herein are large enough to produce the desired effect in the method by which delivery occurs. The dosage
5 should not be so large as to cause adverse side effects, such as unwanted cross-reactions, anaphylactic reactions, and the like. Generally, the dosage will vary with the age, condition, sex and extent of the disease in the subject and can be determined by one of skill in the art. The dosage can be adjusted by the individual physician based on the clinical condition of the subject involved. The dose, schedule of doses
10 and route of administration can be varied.

The efficacy of administration of a particular dose of the compounds or compositions according to the methods described herein can be determined by evaluating the particular aspects of the medical history, signs, symptoms, and objective laboratory tests that are known to be useful in evaluating the status of a
15 subject in need of attention for the treatment of cancer, arthritis, or other diseases and/or conditions. These signs, symptoms, and objective laboratory tests will vary, depending upon the particular disease or condition being treated or prevented, as will be known to any clinician who treats such patients or a researcher conducting experimentation in this field. For example, if, based on a comparison with an
20 appropriate control group and/or knowledge of the normal progression of the disease in the general population or the particular individual: 1) a subject's physical condition is shown to be improved (e.g., a tumor has partially or fully regressed), 2) the progression of the disease or condition is shown to be stabilized, or slowed, or reversed, or 3) the need for other medications for treating the disease or condition is
25 lessened or obviated, then a particular treatment regimen will be considered efficacious.

Any of the compounds represented by Formula I can be used therapeutically in combination with a pharmaceutically acceptable carrier. In another aspect, any of the compounds represented by Formula I can be used prophylactically, *i.e.*, as a preventative agent, with a pharmaceutically acceptable carrier. The compounds
5 described herein can be conveniently formulated into pharmaceutical compositions composed of one or more of the compounds in association with a pharmaceutically acceptable carrier. *See, e.g., Remington's Pharmaceutical Sciences*, latest edition, by E.W. Martin Mack Pub. Co., Easton, PA, which discloses typical carriers and conventional methods of preparing pharmaceutical compositions that can be used in
10 conjunction with the preparation of formulations of the compounds described herein and which is incorporated by reference herein. Such pharmaceutical carriers, most typically, would be standard carriers for administration of compositions to humans and non-humans, including solutions such as sterile water, saline, and buffered solutions at physiological pH. Other compounds will be administered according to
15 standard procedures used by those skilled in the art.

The pharmaceutical compositions described herein can include, but are not limited to, carriers, thickeners, diluents, buffers, preservatives, surface active agents and the like in addition to the molecule of choice. Pharmaceutical compositions can also include one or more additional active ingredients such as antimicrobial agents,
20 antiinflammatory agents, anesthetics, and the like.

The compounds and pharmaceutical compositions described herein can be administered to the subject in a number of ways depending on whether local or systemic treatment is desired, and on the area to be treated. Thus, for example, a compound or pharmaceutical composition described herein can be administered as
25 an ophthalmic solution and/or ointment to the surface of the eye. Moreover, a compound or pharmaceutical composition can be administered to a subject vaginally, rectally, intranasally, orally, by inhalation, or parenterally, for example, by

intradermal, subcutaneous, intramuscular, intraperitoneal, intrarectal, intraarterial, intralymphatic, intravenous, intrathecal and intratracheal routes. Parenteral administration, if used, is generally characterized by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms
5 suitable for solution or suspension in liquid prior to injection, or as emulsions. A more recently revised approach for parenteral administration involves use of a slow release or sustained release system such that a constant dosage is maintained. *See, e.g.,* U.S. Patent No. 3,610,795, which is incorporated by reference herein for its teaching of sustained release systems.

10 Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions which can also contain buffers, diluents and other suitable additives. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water,
15 alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives, such as
20 antimicrobials, anti-oxidants, chelating agents, and inert gases and the like, can also be present.

Formulations for topical administration can include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like can
25 be necessary or desirable.

Compositions for oral administration can include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets, or

tablets. Thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders can be desirable.

EXAMPLES

The following examples are put forth so as to provide those of ordinary skill
5 in the art with a complete disclosure and description of how the compounds, compositions, articles, devices, and/or methods described and claimed herein are made and evaluated, and are intended to be purely exemplary and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.)
10 but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in °C or is at ambient temperature, and pressure is at or near atmospheric. There are numerous variations and combinations of reaction conditions, e.g., component concentrations, desired solvents, solvent mixtures, temperatures, pressures and other reaction ranges and conditions that can
15 be used to optimize the product purity and yield obtained from the described process. Only reasonable and routine experimentation will be required to optimize such process conditions.

Example 1

(*cis*)-2-[[[4-Biphenyl)sulfonyl]amino]cyclohexanecarboxylic acid: A stirred
20 solution of *cis*-2-amino-1-cyclohexanecarboxylic acid (2.4g, 16.76 mmol) and sodium carbonate (3.553g, 33.52 mmol) in dioxane:water (168:84 mL) was cooled in ice-water bath and to the cold solution was added biphenyl-4-sulfonyl chloride (4.85g, 19.15 mmol) in one portion. The reaction mixture was stirred in the cold bath for 2 hours. After allowing the reaction mixture to attain room temperature, the
25 mixture was stirred for an additional 48 hours. The mixture was then poured into 10% aqueous citric acid (500mL) and the mixture was stirred for 2 hours. The solid obtained was collected by filtration. The solid was stirred with 1N aqueous sodium

hydroxide and the solution was filtered to remove any insoluble material. The alkaline aqueous filtrate was cooled and acidified to pH 1 with concentrated aqueous hydrochloric acid. The solid obtained was collected by filtration, washed with water and dried in air to obtain the desired acid as a colorless solid. The procedure yielded
5 3.78g (63%) of (*cis*)-2-[[*(4*-Biphenyl)sulfonyl]amino]cyclohexanecarboxylic acid, which had a melting point of 188–190°C. Mass spectrometry revealed molecular ion peak (MH)⁺ at *m/z* 360. Proton NMR analysis in CDCl₃ resulted in the following chemical shifts (δ): 1.2–2.1 (m, 8H, cyclohexyl methylenes), 2.70–2.85 (m, 1H, C-1 H), 3.40–3.55 (m, 1H, C-2 H), 5.95 (d, 1H, SO₂NH), 7.35–8.0 (m, 9H, biphenyl-H),
10 8.75 (broad s, 1H, CONH), and 10.45 (broad s, 1H, N-OH), where s is a singlet, d is a doublet, m is a multiplet.

Example 2

(*cis*)-*N*-Hydroxy-2-[[*(4*-Biphenyl)sulfonyl]amino]cyclohexanecarboxamide (compound 1c): A solution of the acid obtained in Example 1 (0.80g, 2.23 mmol) in
15 CH₂Cl₂ (30mL) was cooled in an ice bath and treated with oxalyl chloride (1.415g, 11.15 mmol) followed by one drop of *N,N*-dimethylformamide as a catalyst. The reaction mixture was then allowed to warm to room temperature and stir at room temperature for 2 hours. Volatile material from the reaction mixture was then removed under reduced pressure, and the residue was dried under high vacuum for
20 an hour. The acid chloride thus obtained was dissolved in anhydrous tetrahydrofuran (10mL), cooled to 0°C, and treated dropwise with *O*-(trimethylsilyl)hydroxylamine (2.35g, 22.3 mmol). The mixture was allowed to attain room temperature and stir overnight. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in ethyl acetate (200mL). The solution was successively
25 washed with 1 N hydrochloric acid (100mL) and brine (100mL). The organic layer was dried with anhydrous sodium sulfate, filtered, and the solvent was removed under reduced pressure. The solid thus obtained was recrystallized from ethyl

acetate/hexane to yield 0.63g (75%) of the desired product 1c. The product 1c had a melting point of 74–76°C. Mass spectrometry revealed a molecular ion peak (MH)⁺ at *m/z* 375. Proton NMR analysis in DMSO-*d*₆ resulted in the following chemical shifts (δ): 1.0–2.1 (m, 8H, cyclohexyl methylenes), 2.30–2.40 (m, 1H, C-1 H), 3.20–3.35 (m, 1H, C-2 H), 7.35–8.00 (m, 10H, biphenyl-H and SO₂NH), 8.75 (broad s, 1H, CONH), and 10.45 (broad s, 1H, N-OH).

Example 3

(*trans*)-2-[[[4-Biphenyl)sulfonyl]amino]cyclohexanecarboxylic acid: This compound was prepared from *trans*-2-amino-1-cyclohexanecarboxylic acid (1.0g, 7.0 mmol) by reacting it with sodium carbonate (1.48g, 14.0 mmol) and biphenyl-4-sulfonyl chloride (2.02g, 8.0 mmol) in dioxane:water (70:35 mL) in the same manner as described in Example 1. The yield was 1.48g (59%), the melting point was 220–222°C, and mass spectrometry revealed a molecular ion peak (MH)⁺ at *m/z* 360.

Example 4

(*trans*)-*N*-Hydroxy-2-[[[4-Biphenyl)sulfonyl]amino]cyclohexanecarboxamide (compound 2c): A solution of the acid obtained in Example 3 (0.56g, 1.56 mmol) was reacted with oxalyl chloride (0.99g, 7.8 mmol) followed by *O*-(trimethylsilyl)-hydroxylamine (1.64g, 15.6 mmol) as described in Example 2 to obtain 0.23g (40%) of 2c. The melting point was 212–214°C. Mass spectrometry revealed a molecular ion peak (MH)⁺ at *m/z* 375. Proton NMR analysis in DMSO-*d*₆ resulted in the following chemical shifts (δ): 0.90–1.85 (m, 8H, cyclohexyl methylenes), 1.90–2.05 (m, 1H, C-1 H), 3.25–3.45 (m, 1H, C-2 H), 7.40–7.90 (m, 10H, biphenyl-H and SO₂NH), 8.75 (broad s, 1H, CONH), and 10.24 (broad s, 1H, N-OH).

Example 5

(*cis*)-2-[(4-Phenoxybenzenesulfonyl)amino]cyclohexanecarboxylic acid:

This compound was prepared from *cis*-2-amino-1-cyclohexanecarboxylic acid (1.0g, 7.0 mmol) by reacting it with sodium carbonate (1.48g, 14.0 mmol) and 4-
5 phenoxybenzenesulfonyl chloride (2.25g, 8.38 mmol) in dioxane:water (40:20 mL) in the same manner as described in Example 1. This yielded 2.23g (85%). The melting point was 116–118°C. Mass spectrometry revealed a molecular ion peak (MH)⁺ at *m/z* 376. Proton NMR analysis in CDCl₃ resulted in the following chemical shifts (δ): 1.2–2.2 (m, 8H, cyclohexyl methylenes), 2.75–2.85 (m, 1H, C-1
10 H), 3.62–3.50 (m, 1H, C-2 H), 5.92 (d, 1H, SO₂NH), and 6.99–7.86 (m, 9H, aryl-H).

Example 6

(*cis*)-*N*-Hydroxy-2-[(4-Phenoxybenzenesulfonyl)amino]cyclohexanecarbox-

amide (compound 1d): A solution of the acid obtained in Example 5 (0.56g, 1.49 mmol) was reacted with oxalyl chloride (2M solution in CH₂Cl₂, 3.7mL, 7.4 mmol) followed by *O*-(trimethylsilyl)hydroxylamine (1.56g, 14.91 mmol) as described in
15 Example 2 to obtain 0.275g (48%) of the desired product. The melting point was 66–68°C. Mass spectrometry revealed a molecular ion peak (MH)⁺ at *m/z* 391. Proton NMR analysis in DMSO-*d*₆ resulted in the following chemical shifts (δ): 1.06–1.94 (m, 8H, cyclohexyl methylenes), 2.29–2.36 (m, 1H, C-1 H), 3.19–3.28 (m,
20 1H, C-2 H), 7.08–7.86 (m, 10H, biphenyl-H and SO₂NH), 8.8 (broad s, 1H, CONH), and 10.4 (broad s, 1H, N-OH).

Example 7

(*trans*)-2-[(4-Phenoxybenzenesulfonyl)amino]cyclohexanecarboxylic acid:

This compound was prepared from *trans*-2-amino-1-cyclohexanecarboxylic acid
25 (1.0g, 7.0 mmol) by reacting it with sodium carbonate (1.48g, 14.0 mmol) and 4-phenoxybenzenesulfonyl chloride (2.25g, 8.38 mmol) in dioxane:water (40:20 mL)

in the same manner as described in Example 1. This yielded 1.275g (48%). The melting point was 192–194°C. Mass spectrometry revealed a molecular ion (MH)⁺ peak at *m/z* 376.

Example 8

5 (*trans*)-*N*-Hydroxy-2-[(4-Phenoxybenzenesulfonyl)amino]cyclohexane-carboxamide (compound 2d): A solution of the acid obtained in Example 7 (1.00g, 2.66 mmol) was reacted with oxalyl chloride (2M solution in CH₂Cl₂, 6.65mL, 13.3 mmol) followed by *O*-(trimethylsilyl)hydroxylamine (2.8g, 26.64 mmol) as described in Example 2 to obtain 0.48g (46%) of the desired product. The melting
10 point was 182–184°C. Mass spectrometry revealed a molecular ion (MH)⁺ peak at *m/z* 391. Proton NMR analysis in CDCl₃ resulted in the following chemical shifts (δ): 0.98–1.96 (m, 8H, cyclohexyl methylenes), 2.05–2.16 (m, 1H, C-1 H), 3.50–3.67 (broad hump, 1H, C-2 H), 6.15 (broad hump, 1H, SO₂NH), 7.01–7.88 (m, 9H, aryl-H), and 8.98 (broad s, 1H, CONH).

15 Example 9

 (*cis*)-2-[[[4-Phenylazo)benzenesulfonyl]amino]cyclohexanecarboxylic acid: This compound was prepared from *cis*-2-amino-1-cyclohexanecarboxylic acid (1.0g, 7.0 mmol) by reacting it with sodium carbonate (1.48g, 14.0 mmol) and 4-(phenylazo)benzenesulfonyl chloride (2.35g, 8.38 mmol) in dioxane:water (40:20
20 mL) in the same manner as described in Example 1. This yielded 0.915g (33%). Mass spectrometry revealed a molecular ion peak (MH)⁺ at *m/z* 388. Proton NMR analysis in CDCl₃ resulted in the following chemical shifts (δ): 1.2–1.9 (m, 8H, cyclohexyl methylenes), 1.98–2.14 (m, 1H, C-1 H), 3.45–3.60 (m, 1H, C-2 H), 5.9 (broad hump, 1H, SO₂NH), and 7.50–8.05 (m, 9H, aryl-H).

Example 10

(*cis*)-*N*-Hydroxy-2-[[[(4-Phenylazo)benzenesulfonyl]amino]cyclohexane-carboxamide (compound 1g): A solution of the acid obtained in Example 9 (0.915g, 2.36 mmol) was reacted with oxalyl chloride (2M solution in CH₂Cl₂, 5.91mL, 11.82 mmol) followed by *O*-(trimethylsilyl)hydroxylamine (2.48g, 23.64 mmol) as described in Example 2 to obtain 0.44g (46%) of the desired product. The melting point was 162–164°C. Mass spectrometry revealed a molecular ion peak (MH)⁺ at *m/z* 403. Proton NMR analysis in CDCl₃ resulted in the following chemical shifts (δ): 0.71–2.05 (m, 8H, cyclohexyl methylenes), 2.6–2.7 (m, 1H, C-1 H), 2.30–2.41 (m, 1H, C-2 H), 6.1–6.2 (broad hump, SO₂NH), 7.5–8.1 (m, 10 H, aryl-H), and 8.3–8.6 (broad hump, 2H, CONH, N-OH).

Example 11

(*trans*)-2-[[[(4-Phenylazo)benzenesulfonyl]amino]cyclohexanecarboxylic acid. This compound was prepared from *trans*-2-amino-1-cyclohexanecarboxylic acid (1.0g, 7.0 mmol) by reacting it with sodium carbonate (1.48g, 14.0 mmol) and 4-(phenylazo)benzenesulfonyl chloride (2.35g, 8.38 mmol) in dioxane:water (40:20 mL) in the same manner as described in Example 1. This yielded 1.02g (37%). The melting point was 246–248°C. Mass spectrometry revealed a molecular ion peak at (MH)⁺ *m/z* 388. Proton NMR analysis in CDCl₃ resulted in the following chemical shifts (δ): 1.0–1.98 (m, 8H, cyclohexyl methylenes), 2.05–2.14 (m, 1H, C-1 H), 3.35–3.45 (m, 1H, C-2 H), 4.88–4.99 (broad hump, 1H, SO₂NH), and 7.5–8.1 (m, 9H, aryl-H).

Example 12

(*trans*)-*N*-Hydroxy-2-[[[(4-Phenylazo)benzenesulfonyl]amino]cyclohexane-carboxamide (compound 2g): A solution of the acid obtained in Example 11 (1.00g,

2.58 mmol) was reacted with oxalyl chloride (2M solution in CH_2Cl_2 , 6.45mL, 12.9 mmol) followed by *O*-(trimethylsilyl)hydroxylamine (2.72g, 25.84 mmol) as described in Example 2 to obtain 0.35g (33%) of the desired product. The melting point was 194–196°C. Mass spectrometry revealed a molecular ion peak $(\text{MH})^+$ at *m/z* 403. Proton NMR analysis in CDCl_3 resulted in the following chemical shifts (δ): 0.98–1.98 (m, 8H, cyclohexyl methylenes), 2.10–2.22 (m, 1H, C-1 H), 3.60–3.78 (m, 1H, C-2 H), 6.35 (d, 1H, SO_2NH), 7.50–8.15 (m, 9H, aryl-H), and 8.9–9.0 (broad hump, 2H, CONH, N-OH).

Example 13

Enzyme Inhibition Assays: The enzyme inhibition kinetics (IC_{50}) of nonpeptidyl compounds 1c and 2c were determined using purified MMP-2, MMP-3, and MMP-9 using standard fluorometric substrate-degradation assays. Human rMMP-2 and -9 were purified in active forms, and purified MMP-1, -3 and -13 were obtained from commercial sources (Chemicon; Temecula, CA) in zymogen form and were activated by 1mM APMA or PCMB treatment (2 hrs at 37 °C). For kinetic studies, a standard fluorimetric assay based on hydrolysis of fluorogenic synthetic substrate (1 μM of McaPLGLDpaAR) was used. For each assay, target MMP (1 μM) and increasing concentrations of test compound were incubated at 25°C for 30-60 min and the rate of substrate hydrolysis was measured by Perkin-Elmer fluorometer at an excitation 328nm and an emission 393nm settings. Both concentration and time-dependent cleavage were monitored. Controls included trypsin (non-specific cleavage) and bacterial collagenase (total cleavage) and other zinc-protease (ACE, endopeptidase). IC_{50} values (the concentration at which 50% enzyme activity is inhibited) were determined by plots of % activity versus negative log of agent concentration. The IC_{50} values were converted to K_i values using the equation, $K_i = \text{IC}_{50} / (1 + S/K_m)$. The K_m values (μM) were calculated from the K_i values of 3-5 separate experiments. Inhibition kinetics of various MMPs were also determined

using substrate-degradation ELISA kits from Chemicon (Temecula, CA). See C. Knight, *et al.*, *FEBS Lett.*, 1992, 296, 263-266; L. Windsor, *et al.*, *Biochem. Biophys. Acta* 1977, 1334, 261-272, which are incorporated by reference herein for their teachings of enzyme inhibition assays.

5 MMP inhibitor selectivity determinations: Both quantitative labeled-substrate and FITC-biopeptide (matrix-specific) assays were used to examine the selective inhibition of MMP activities. Substrate specificity for MMP-1, -2, and -9 was assessed by degradation of labeled-collagen I and IV. Typically, MMP-dependent biomatrix degradation was monitored in the presence and absence of
10 candidate compounds (0.5-500 μ m) after 60-90min incubations. Inhibitory potency and selectivity was also analyzed using a gelatin or collagen zymography followed by densitometric analysis. Labeled collagen I or IV degradation assay (zymography) was used for MMP-1, -2 and -9 activities and their inhibition. Bacterial collagenase and EDTA (10mM) was included as positive and negative controls. *In situ*
15 zymography film technique was also used for measurement of MMP-2 and MMP-9 (gelatinolytic) activities and their inhibition by drug treatment in target cells and tissue samples. For evaluation of MMP selectivity, two human HT-1080 cancer cell sublines, which produce in abundance different levels of MMP-2 and MMP-9 were used. The HT-1080-B subline produces predominantly MMP-9 (2mg/L) while HT-
20 1080-R cells are abundant in MMP-2 (1mg/L). Both cell lines also secrete small amounts of other MMPs including MMP-1 and MMP-3 (< 1%). See G. Siegal, *et al.*, *Cancer Lett.* 1993, 69, 123-132; L. Goodly, *et al.*, *Tumor Biology* 1994, 15, 326-336; M. Ikeda, *et al.*, *Clin. Cancer. Res.* 2000, 6, 3290-3296.

The results presented in Table 1 demonstrate that both sulfonamide
25 compounds selectively inhibit MMP-2 and MMP-9 activities, over MMP-3 and endopeptidase. These inhibition profiles were compared with a potent broad range MMP inhibitor GM-6001, which is available from Chemicon (Temecula, CA).

Negative controls included other proteases, *e.g.*, trypsin or amidopeptidase, whose activities were unaffected (up to 0.1mM).

Table 1: Inhibition Potencies of Nonpeptidyl Compounds Against Selected Enzymes.

	Inhibition Potencies, IC ₅₀ (nM)			
Compound	MMP-2	MMP-9	MMP-3 ^a	Endopeptidase
2c	125	145	>3000	>20000
1c	150	175	>3000	>20000
GM-6001	2	2	80	not detectable

5 ^a Minienzyme used.

Example 14

Tumor Invasion and Angiogenesis Assays: The multi-step metastatic cascade involves MMP-mediated tissue matrix degradation, angiogenesis, tumor cell migration/invasion and subsequent colonization at distant sites (M. Crocket, *et al.*,
10 *Biochem Soc. Symp.*, 1998, 63, 295-313; D. Keiner, *et al.*, *Metastasis Rev.*, 1990, 9, 289-303; J. MacDougall, *et al.*, *Mol. Med. Today*, 2000, 64, 149-56). These functional events have been mostly studied using biomatrix co-culture systems such as collagen gels and MATRIGELTM (available from Becton-Dickinson, Bedford, Mass; see R. Auerbach, *et al.*, *Pharm. Ther.*, 1991, 51, 1-11; H. Kleinman, *et al.*,
15 *Biochem.*, 1986, 25, 312-318). However, these models exhibit compositional complexity and biological limitations. For example, the mouse tumor derived MATRIGELTM contains major mitogenic and differentiation factors as well as proteases which may trigger undefined cell-matrix interactions (S. Vukicevik, *et al.*,

Exp. Cell Res., 1992, 202, 1-8). To overcome these difficulties a new human biomatrix, Amgel, which is free of collagenases and mitogenic factors, was used for examining the functional behavior of target cells (G. Siegel, *et al.*, *Cancer Lett.*, 1993, 69, 123-132).

5 The hallmarks of the Amgel system are that it mimics a physiological matrix barrier, and Amgel alone is neither angiogenic nor tumorigenic, but exhibits controlled bioactivity both *in vitro* and *in vivo*. Also, the Amgel bioassay can identify single modulators (natural and synthetic agents) of human cell invasion, motility and angiogenesis. Moreover, the Amgel biomatrix contains both human
10 collagen I and IV, the *in vivo* footprints of MMP-1 activity and MMPs-2 and -9 activities, respectively. Thus, Amgel bioassays are useful for identifying the selectivity of MMP inhibitors while discriminating their effects on different stages of tumorigenesis. Accordingly, to validate if MMP inhibition translated into a biological effect, Amgel human tumor invasion and angiogenesis models were
15 employed.

Human tumor cell invasion bioassay: Amgel-coated filters (8 μ m) were used as tissue biomatrix barriers placed between Lucite chambers. Labeled cells (50,000) were seeded onto reconstituted-Amgel filters (75 μ g) and the lower chambers were filled with media containing 5% dialyzed serum. Test compounds (0.5-100 μ M)
20 were added and incubated for 72 hours. Contents from the lower chamber were collected to resolve cell-bound and free radioactivity. See G. Siegal, *et al.*, *Cancer Lett.* 1993, 69, 123-132; L. Goodly, *et al.*, *Tumor Biology* 1994, 15, 326-336; M. Ikeda, *et al.*, *Clin. Cancer. Res.* 2000, 6, 3290-3296; R. Singh, *et al.*, *In Vitro Cell Develop. Biol.*, 2002, 38, 11.

25 Human tumor angiogenesis bioassay: Angiogenesis was induced by coculture of human endothelial cells (HUVECs) with tumor cells or defined angiogenic factor (FGF). Tumor cells media (from 24-hour serum-free cultures)

were used to induce a biological angiogenic response using human glioma cell lines producing different concentrations of VEGF or by using purified factors such as 20 μ M FGF. Endothelial cells were seeded onto Amgel-coated filters and induced with a angiogenic media +/- candidate MMP inhibitors. Varying concentrations of test compounds (0.5-500 μ M) and incubation times (1-7 days) were applied, and endothelial cell differentiation (sprouting and tubule-like capillary formation) were examined by light microscopy and a computerized digital system.

Evaluation of the compounds 1c and 2c in Amgel human tumor invasion and angiogenesis models (72 hours incubation at 50-100 μ M) was performed using two human HT-1080 cancer cell sublines (HT-1080-B and HT-1080-R). The results of these assays with compound 2c are shown in Table 2 and Figure 1 (data for compound 1c not shown).

Table 2: Inhibition Potencies of Compound 2c in Cell-Function Assays.

Cell line	% Inhibition	
	Tumor invasion	Tumor angiogenesis
HT-1080R	60 \pm 5	40 \pm 6
HT-1080B	50 \pm 8	45 \pm 4
HF	No effect	No effect

Functional evaluation of compounds 1c and 2c revealed a marked (40-60%) reduction in the invasiveness and angiogenesis of highly tumorigenic human HT-1080 cell lines. Also, neither compound 1c nor compound 2c had any effect on nontumorigenic human fibroblasts cells (HF). Further, compounds 1c and 2c were

observed to have no effect on the cell proliferation and cell viability at the tested concentrations (data not shown).


5 The arrest of tumor invasion by compounds 1c and 2c paralleled their profiles in inhibiting MMP-2 and MMP-9 activities, as measured by degree of gelatin degradation by electrophoresed cell media. The inhibition potencies of the compounds in the parental HT-1080 cells producing both MMP-2 and MMP-9 were much higher (> 70%) than their inhibition potencies in cell lines producing a single form of MMP. These observations indicate that compounds 1c and 2c produce synergistic effects through simultaneous inhibition of MMP-2 and MMP-9.

10 Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the compounds, compositions and methods described herein.

15 Various modifications and variations can be made to the compounds, compositions and methods described herein. Other aspects of the compounds, compositions and methods described herein will be apparent from consideration of the specification and practice of the compounds, compositions and methods disclosed herein. It is intended that the specification and examples be considered as exemplary.

What is claimed is:

1. A compound having the following formula:



wherein



X is $(\text{CH}_2)_n\text{O}$, $(\text{CH}_2)_n\text{S}$, $(\text{CH}_2)_n\text{NR}^1$, $(\text{CH}_2)_n(\text{CH}_2)$, or $\text{CH}=\text{CH}$, wherein $n = 0, 1$, or 2 ;

R and R^1 are, independently, a substituted or unsubstituted alkyl, alkenyl, alkynyl, aryl, heteroaryl group, cycloalkyl, heterocycloalkyl, cycloalkenyl, or heterocycloalkenyl; and


Z is NH or CH_2 ;

or a pharmaceutically acceptable salt thereof.


2. The compound of claim 1, wherein Z is NH.
3. The compound of claim 1, wherein Z is CH_2 .
4. The compound of claims 2 or 3, wherein R is a substituted or unsubstituted aryl or heteroaryl group.
5. The compound of claims 2 or 3, wherein R is a substituted aryl group of the following formula:



6. The compound of claims 2 or 3, wherein R is:




7. The compound of claims 2 or 3, wherein X is $(CH_2)_n(CH_2)$ and $n = 1$.
 8. The compound of claims 2 or 3, wherein X is $CH=CH$.
 9. The compound of claim 1, wherein the compound is




or a pharmaceutically acceptable salt thereof.

10. The compound of claim 1, wherein the compound is




or a pharmaceutically acceptable salt thereof.

11. The compound of claim 1, wherein the compound is




or a pharmaceutically acceptable salt thereof.

12. The compound of claim 1, wherein the compound is



or a pharmaceutically acceptable salt thereof.

13. The compound of claim 1, wherein the compound is



or a pharmaceutically acceptable salt thereof.

14. The compound of claim 1, wherein the compound is a selective modulator of a MMP.
15. The compound of claim 1, wherein the compound is a modulator of human tumor metastasis.
16. The compound of claim 1, wherein the compound is a modulator of MMP-2, MMP-9, or a mixture thereof, *in vitro*.
17. The compound of claim 1, wherein the compound is a selective inhibitor of a MMP.
18. The compound of claim 1, wherein the compound is an inhibitor of human tumor metastasis.
19. The compound of claim 1, wherein the compound is an inhibitor of MMP-2, MMP-9, or a mixture thereof, *in vitro*.
20. A pharmaceutical composition, comprising the compound of claim 1 and a pharmaceutical carrier.
21. The composition of claim 20, wherein the compound is the compound of claims 9 or 10.

22. The composition of claim 20, further comprising an anti-cancer agent.
23. A method for using the compound of claim 1, comprising:


administering an amount effective for modulation of a MMP of at least one compound of claim 1 to an environment comprising the MMP.
24. The method of claim 23, wherein the MMP is MMP-2, MMP-9, or a mixture thereof.
25. The method of claim 23, wherein the at least one compound is the compound of claim 9.
26. The method of claim 23, wherein the at least one compound is the compound of claim 10.
27. The method of claim 23, wherein the amount effective for modulation is equivalent to an amount effective for inhibition.
28. The method of claim 27, wherein inhibition is characterized by an IC_{50} less than about 3000nM.
29. The method of claim 27, wherein inhibition is characterized by an IC_{50} less than about 200nM.
30. A method for using the compound of claim 1, comprising:

administering an amount effective for modulation of tumor metastasis of at least one compound of claim 1 to a cell.
31. The method of claim 30, wherein the amount effective for modulation is equivalent to the amount effective for inhibition.
32. The method of claim 30, wherein the cell is a HT-1080 cell.

33. The method of claim 30, wherein inhibition is measured by arrest of tumor invasion.
34. The method of claim 30, wherein inhibition is measured by arrest of tumor angiogenesis.
35. A method for treating a subject with cancer comprising administering an effective amount of the compound of claim 1 to a subject in need of the treatment.
36. The method of claim 35, wherein the cancer is a carcinoma, melanoma, leukemia, or adenoma.
37. The method of claim 35, wherein the compound of claim 1 is part of an anti-cancer cocktail.
38. The method of claim 35, wherein the subject is a human.
39. A method for preventing cancer in a subject comprising administering an effective amount of the compound of claim 1 to a subject.
40. A method for treating a subject with arthritis comprising administering an effective amount of the compound of claim 1 to a subject in need of the treatment.

ABSTRACT

Disclosed are selective inhibitors of matrix metalloproteinases represented by the following formula (I).



(I)

wherein X is $(\text{CH}_2)_n\text{O}$, $(\text{CH}_2)_n\text{S}$, $(\text{CH}_2)_n\text{NR}^1$, $(\text{CH}_2)_n(\text{CH}_2)$, or $\text{CH}=\text{CH}$, wherein $n = 0, 1$, or 2 ; R and R^1 are, independently, a substituted or unsubstituted alkyl, alkenyl, alkynyl, aryl, heteroaryl group, cycloalkyl, heterocycloalkyl, cycloalkenyl, or heterocycloalkenyl; and Z is NH or CH_2 ; or a pharmaceutically acceptable salt thereof. Also disclosed are methods of making such compounds and methods of using such compounds to inhibit tumor progression and to treat diseases such as arthritis.

Attorney Docket No.: 19044.0057UI
Inventor: Subramaniam Ananthan
Filed: March 22, 2004
Figure 1 of 1




Fig. 1

From the INTERNATIONAL BUREAU

PCTNOTIFICATION CONCERNING
SUBMISSION OR TRANSMITTAL
OF PRIORITY DOCUMENT

(PCT Administrative Instructions, Section 411)

To:

KATZ, Mitchell, A.
Needle & Rosenberg, P.C.
Suite 1000
999 Peachtree Street
Atlanta, GA 30309-3915
ETATS-UNIS D'AMERIQUEDate of mailing (day/month/year)
24 May 2005 (24.05.2005)Applicant's or agent's file reference
19044.0057P1International application No.
PCT/US05/009263

International publication date (day/month/year)

IMPORTANT NOTIFICATIONInternational filing date (day/month/year)
21 March 2005 (21.03.2005)Priority date (day/month/year)
22 March 2004 (22.03.2004)

Applicant

SOUTHERN RESEARCH INSTITUTE et al

1. By means of this Form, which replaces any previously issued notification concerning submission or transmittal of priority documents, the applicant is hereby notified of the date of receipt by the International Bureau of the priority document(s) relating to all earlier application(s) whose priority is claimed. Unless otherwise indicated by the letters "NR", in the right-hand column or by an asterisk appearing next to a date of receipt, the priority document concerned was submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b).
2. (If applicable) The letters "NR" appearing in the right-hand column denote a priority document which, on the date of mailing of this Form, had not yet been received by the International Bureau under Rule 17.1(a) or (b). Where, under Rule 17.1(a), the priority document must be submitted by the applicant to the receiving Office or the International Bureau, but the applicant fails to submit the priority document within the applicable time limit under that Rule, the attention of the applicant is directed to Rule 17.1(c) which provides that no designated Office may disregard the priority claim concerned before giving the applicant an opportunity, upon entry into the national phase, to furnish the priority document within a time limit which is reasonable under the circumstances.
3. (If applicable) An asterisk (*) appearing next to a date of receipt, in the right-hand column, denotes a priority document submitted or transmitted to the International Bureau but not in compliance with Rule 17.1(a) or (b) (the priority document was received after the time limit prescribed in Rule 17.1(a) or the request to prepare and transmit the priority document was submitted to the receiving Office after the applicable time limit under Rule 17.1(b)). Even though the priority document was not furnished in compliance with Rule 17.1(a) or (b), the International Bureau will nevertheless transmit a copy of the document to the designated Offices, for their consideration. In case such a copy is not accepted by the designated Office as the priority document, Rule 17.1(c) provides that no designated Office may disregard the priority claim concerned before giving the applicant an opportunity, upon entry into the national phase, to furnish the priority document within a time limit which is reasonable under the circumstances.

<u>Priority date</u>	<u>Priority application No.</u>	<u>Country or regional Office or PCT receiving Office</u>	<u>Date of receipt of priority document</u>
22 March 2004 (22.03.2004)	60/555,380	US	25 April 2005 (25.04.2005)

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No. +41 22 740 14 35

Authorized officer

Boccard Paulette

Facsimile No. +41 22 338 87 40
Telephone No. +41 22 338 8147